

Summary statistics for continuous data

Tabriz University of Medical
Sciences
Standard Workshop on
Systematic Reviews _ October
2012

Dr. Shayesteh Jahanfar,
University of British Columbia

Outline

- identify continuous outcomes
- understand how to summarise continuous data and pool studies with:
 - measures on the same scale
 - measures on different scales
- recognise some of the challenges of continuous data



Types of data

- Binary data
- Counts of infrequent events (e.g. number of strokes)
- Short ordinal scales (e.g. pain grades: none/mild/moderate/severe)
- Long ordinal scales (e.g. disability scales)
- Continuous data (e.g. blood pressure)
- Censored data (e.g. survival times)

What are continuous data?

- data with an infinite number of values that are equally spaced
- example: height - it can be measured along a numerical continuum of centimetres, metres or inches, feet
 - a person can be 175.24678cm tall, assuming the measurement instrument is accurate enough
 - the difference between 160 and 161cm, and 180 and 181cm, is the same



Long ordinal scales

- sometimes treated as continuous data
- but not true continuous because
 - they have a finite number of distinct values
 - there are gaps in the continuum
- have multiple, ordered categories which imply magnitude
 - e.g. one category is greater or lesser than another
- spacing between categories is not numerically equivalent
- approach 'continuous' with increasing categories



What continuous data can we combine?

- data represent continuous measures
- the mean value is in the middle (distribution is roughly symmetrical)
- measurements are made on all participants (not censored or survival type data)
- data are available for both groups in each trial



What data is needed?

	Mean	SD	Sample size
Treatment	m_t	sd_t	n_t
Control	m_c	sd_c	n_c

Meta-analysis of continuous data

- calculate a single summary statistic to represent the effect found in each study
- Summary statistics combined in meta-analysis
- 2 options
 - mean difference
 - standardised mean difference



Mean difference


- outcomes measured in same unit using same scale (e.g. blood pressure as mmHg)
- pooled analysis in “natural units” and therefore easy to interpret
- studies weighted according to the inverse of the variance (a function of size and SD)

MD = mean on treatment – mean on control



Mean difference: example

Review: Caffeine for daytime 'sluggishness'. (version with data)
 Comparison: 01 Caffeinated Coffee versus Decaffeinated Coffee
 Outcome: 03 Irritability at 30 minutes - INAS scale (1-50, high score worse)



Standardised mean difference


- Outcome is same concept measured on different scales, the values must be transformed to a common scale before pooling
- Sometimes scale factors are known and transformations are made directly (e.g weight)
- Standardised mean difference calculated as:

$$\frac{\text{Difference in means between groups}}{\text{Average standard deviation}}$$




Standardised mean difference

Beck Irritability Scale (1-30)



Irritability Negativity Affectivity Subscale (1-50)



Different scales but averages mean the same thing
(i.e. average person is just as irritable!)

Measurements on different scales

Comparing irritability at 30 minutes between caffeinated coffee and decaffeinated coffee


Trial	Caffeinated N. mean (SD)	Decaffeinated N. mean (SD)	Irritability scale
Moccona 1998	15 23.0 (15.1)	17 31.0 (15.2)	INAS
Nescafe 1998	68 19.0 (15.5)	64 36.0 (17.3)	INAS
Piazza D'oro 2003	35 21.0 (3.2)	37 10.0 (4.20)	BII

High scores on the Beck Irritability Scale (BII) (1-30) good outcomes, while high scores on the Irritability Negative Affectivity Subscale (INAS) (1-50) are poor outcomes



SMD: example

Review: Caffeine for daytime 'sluggishness'. (version with data)
 Comparison: 01 Caffeinated Coffee versus Decaf
 Outcome: 06 Irritability at 30 minutes



RevMan exercise



Change vs endpoint scores

Start of
study

End of study

Treatment group

Score T0

Change in score T

Score T1

Difference in mean
change scores

Difference
in mean
end point
scores

Control group

Score C0

Change in score C

Score C1



Problems with MD and SMD

- what constitutes a clinically important change?
- restrictive eligibility criteria results in smaller standard deviations; therefore these trials given more weight
- mean difference
 - measurements on the same scale are not always comparable (e.g. health care costs in different places, process of care measures)
- standardised mean difference
 - difficult to interpret outcomes in units of SD, but can transform back to units of the scale
 - estimates of variation may not always be comparable making the SD a poor scaling factor



Take home message

- pooling continuous data – use mean difference or standardised mean difference
- check data for skewness
- can calculate SDs from other statistics
- can use either endpoint or change scores

